

Experience of the practical implementation of the Hardware-Software complex "Sensitiv Imago" for diagnostics and treatment of different diseases

At the moment the methods of people's state of health screening evaluation have a wide implementation as pre-clinical links of primary medical care. The most popular of them are methods of electropuncture diagnostics after Voll , diagnostics after Nakatani , termoalgotometry method and other methods of bioresonant diagnostics and therapy [4,5], and the most effective method with device "**Sensitiv Imago**".

In our Center in the period of September 2004 to December 2004 a research work had been carried out (246 patients at the age of 19 to 83 years (123 men and 123 women) were examined) to evaluate the precision and efficiency of the offered method of human organism testing with the Hardware-Software Complex "**Sensitiv Imago**" [1,2,3]; the theoretic bases of method of "**Sensitiv Imago**" correspond to the main principles of bioresonance diagnostics and therapy. To estimate the correspondence of the detected pathology to the clinical diagnoses, some labor, biochemical, cardiologic, functional, roentgenologic and ultrasound investigation

had been done. The therapy was instituted corresponding to the detected diseases.

In device "**Sensitiv Imago**" health improving means (Dietary Supplements) were selected with the help of computer function "Vegeto-test" to select preparations. In some cases the energy information correction of separate organs using the Mor-function had been done , the bioresonant preparations manufactured using the function "Reprinter" were recommended. Besides this the pharmacology preparations were prescribed if indicated. The results of computer test and also the opinions of the patients are given below.

| Diagnoses | Group | Fixed coincidence rate of detected diagnoses with clinical ones, % | Subjective opinions of patients about the therapy and diagnostics, % | Comments |
|--------------------------|-------|--|--|---|
| Vegeto-vascular dystonia | Men | 91% | fair 10 good 70 excellent 20 | Often have the VVD patients who have the EH and on the contrary |
| Essential hypertension | Men | 44% | good 85 excellent 15 | Often have the EH patients who have the VVD and on the contrary |
| Acute bronchitis | Men | 91% | Good 50 excellent 50 | Often can be detected in case of chronic |

| | | | | |
|--------------------------------|-----|------|-------------------------|---|
| | | | | bronchitis |
| Chronic bronchitis | Men | 90% | good 20 excellent 80 | The lung tbs often can be detected which wasn't confirmed by bronchoscopy, fluorography, roentgenography of the lungs |
| Biliary dyskinesia | Men | 93% | fair 22 good 78 | |
| Colon dysbacteriosis | Men | 90% | Good 100 | |
| Otitis | Men | 100% | Good 30 excellent 70 | |
| Deforming osteoarthritis | Men | 82% | Fair 34 good 66 | Often can be detected if any joint pathology is present |
| Intervertebral osteochondrosis | Men | 86% | fair 20 good 80 | |
| Ischemic cardiac disease | Men | 89% | Good 100 | |
| Peptic ulcer of duodenum | Men | 84% | Good 50 excellent 50 | In any cases can't be confirmed by FHDS |
| Gastroduodenitis | Men | 95% | Good 100 | |
| Pyelonephritis | Men | 87% | Good 50 excellent 50 | |
| Urolithiasis | Men | 89% | fair 30 good 70 | In any cases can't be confirmed by US investigation |
| Chronic cholecystitis | Men | 84% | Good 100 | |
| Chronic persistent hepatitis | Men | 84% | fair 20 good 80 | When examined additionally often can be detected an asymptomatic virus infectioncarrier state |
| Insulinindependentdiabetes | Men | 90% | Good 85 Fair 15 | In the most cases one can see the reducing of glucose tolerance when making probe with sugar load |
| Progressing diabetes mellitus | Men | 60% | Good 50 fair 50 | In many cases can't be confirmed by sugar blood probe |

| | | | | |
|-------------------------|-----|-------|--|---|
| Calculous cholecystitis | Men | 65% | Good 55 fair 45 | In a number of cases can't be confirmed by US investigation |
| Total average: | Men | 83,9% | Excellent 17,8 Good 69,4 Fair 12,8 | |

| Diagnoses | Group | Fixed coincidence rate of detected diagnoses with clinical ones, % | Subjective opinions of patients about the therapy and diagnostics, % | Comments |
|--------------------------------|--------------|---|---|---|
| Vegeto-vascular dystonia | women | 93% | Fair 20 Good 80 | the VVD often can be detected at patients who have the EH and on the contrary |
| Essential hypertension | women | 87% | Good 100 | The EH often can be detected at patients who have the VVD and on the contrary |
| Acute bronchitis | women | 89% | Good 50 excellent 50 | Often can be detected in case of chronic bronchitis |
| Chronic bronchitis | women | 87% | good 80 excellent 20 | Often the tbs is detected which wasn't confirmed by bronchoscopy, FG, RG graphy |
| Biliary dyskinesia | women | 95% | Excellent 100 | |
| Colon dysbacteriosis | women | 92% | Good 50 Excellent 50 | |
| Otitis | women | 87% | good 100 | |
| Intervertebral osteochondrosis | women | 84% | Good 40 Excellent 60 | |
| Ischemic cardiac disease | women | 84% | Good 100 | |
| Peptic ulcer of duodenum | women | 84% | Good 65 excellent 35 | In a number of cases can't be confirmed by FGDS |
| Gastroduodenitis | women | 90% | good 70 excellent 30 | |
| Pyelonephritis | women | 87% | good 50 | |

| | | | | |
|-------------------------------|-------|--------|---|--|
| | | | excellent 50 | |
| Urolithiasis | women | 80% | Fair 20 good 80 | In a number of cases can't be confirmed by US investigation |
| Chronic cholecystitis | women | 97% | Excellent 100 | |
| Chronic persistent hepatitis | women | 86% | good 40 excellent 60 | When being additionally examined often can be detected an asymptomatic virus carrier state |
| Insulinindependentdiabetes | women | 95% | good 40 excellent 60 | In the most cases one can see a reduced glucose tolerance when a probe is made with sugar load |
| Progressing diabetes mellitus | women | 70% | fair 50 good 50 | Many cases couldn't be confirmed by sugar blood probe |
| Deforming osteoarthrosis | women | 85% | good 90 excellent 10 | Often can be detected when having any joint pathology |
| Calculous cholecystitis | women | 83% | good 100 | In a number of cases can be confirmed by US investigation |
| Total average: | | 87, 4% | Excellent 32,8 Good 62,4 Fair 4,8 | |

Notes:

- 1.The formulation "Essential hypertension" contained in the software means that any class of III-stage EH can be present.
- 2.The formulation "Insulin-independent diabetes" contained in the software means diabetes mellitus of type II.
- 3.The formulation "Progressing diabetes mellitus" contained in the software means diabetes mellitus of type I.
4. The formulaion " Calculous cholecystitis " contained in the software means cholelithiasis.

Discussing of results

The data presented in the table show that the women's testing is more successful than the men's one. So, the coincidence rate of test results with the clinical diagnoses in the group of women is 87,4% but in the group of men this value is

83,9%. Besides this, in the group of women one can see the tendency to give positive comments to the test and treatment results with using the functions of the software-hardware complex represented in this article. The comparing of the above mentioned results is shown as following diagrams:

Рисунок 1.




FIG.1

Where:

- blue colored part is the histogram showing the coincidence rate of computer test results and of clinical diagnoses in the group of men
- wine-colored column shows the coincidence rate of computer test results and of clinical diagnoses in the group of women




FIG.2

Where:

- - blue colored part is a segment showing the rate of opinions “excellent” in the group of men
- - wine-colored part is a segment showing the rate of opinions “good” in the group of men
- - yellow colored part is a segment showing the rate of opinions “fair” in the group of men



FIG.3

Where:

- blue colored part is a segment showing the rate of opinions “excellent” in the group of women
- - wine colored part is a segment, showing the rate of opinions “good” in the group of women
- yellow colored part is a segment showing the rate of opinions “fair” in the group of women.

Hence, it is visible that the best results of organism test and the results of different health improving influences with using of additional function as informational energy correcting of separate organs and tissues ("Mora-Therapy"), the results of taking the bioresonance preparations fabricated with the help of informational energy transfer function ("Reprinter") have been achieved in the group of women in comparison with the group of men, because the women's organism has some reasons to be considered as an "open" biological system and because the reasons of diseases almost always are the deep changes of energy streams; and at the bioresonance therapy we most have deal with the electromagnetic oscillations, with information streams, which can be characterized as data exchange. The predominance of positive subjective opinions in the group of women can be partly stipulated by the predominance of psychosomatic diseases [6, 7], and by peculiarities of women's psychoemotional status, which can be corrected with bioresonance preparations ("Mora-therapy"); the process of influention can be seen on the monitor. It is also visible that the diseases of gastrointestinal tract and of the hepatobiliary system are more detectable both in the group of men and in the group of women; this fact is also confirmed by other investigation described by other authors [6, 7, 8, 9, 10].

Conclusions:

- The results of computer test of the whole organism with using of the HSC "**Sensitiv Imago**" in the most number of cases (83,9% and 87.4% in the group of men and of women correspondingly) coincide with the clinical diagnoses and with the results obtained with other methods.
- The detectability of diseases in the group of women is some more than in the group of men.
- The diseases of gastrointestinal tract and of the hepatobiliary system are more detectable than other ones.
- In the group of women the positive opinions about the testing with the HSC "**Sensitiv Imago**" and about health improving measures were expressed more often, what is connected with presented psychosomatic diseases and with the features of women's psychoemotional status.

Literature:

1. Гомеопатия и энергоинформатика. Лупичев Н.Л./Москва, «РОЙ», 1994
2. Метод вариационной термоалгометрии в традиционной диагностике. Методические рекомендации МЗ РФ № 99/95, Москва, 2000
Демин С.А., Демина И.Ф., Терехов Е.А., Беляев Е.А., Шарипова М.М
3. ЯТРОГЕННЫЕ ПОРАЖЕНИЯ ГЕПАТОБИЛИАРНОЙ ЗОНЫ И ИХ КОРРЕКЦИЯ. Полунина Т.Е. //Экспериментальная и клиническая

гастроэнтерология, N1, 2003.

4. ПОКАЗАТЕЛИ СЕКРЕЦИИ ХОЛЕЦИСТОКИНИНА, ВЕГЕТАТИВНОЙ РЕГУЛЯЦИИ СЕРДЕЧНОГО РИТМА И УРОВНЯ ТРЕВОЖНОСТИ У БОЛЬНЫХ С МОТОРНОЙ ДИСФУНКЦИЕЙ ЖЕЛЧНОГО ПУЗЫРЯ

Федоров Н.Е., Немцов Л.Н., Солодков А.П., Кичигина Т.Н., Дроздова М.С.

//Экспериментальная и клиническая гастроэнтерология, N1, 2003.

5. РОЛЬ ДИСБИОТИЧЕСКИХ НАРУШЕНИЙ В ЭТИОЛОГИИ И ПАТОГЕНЕЗЕ СИНДРОМА РАЗДРАЖЕННОГО КИШЕЧНИКА

Ручкина И.Н., Парфенов А.И., Осипов Г.А. //Экспериментальная и клиническая гастроэнтерология, N1, 2003.

6. СРАВНИТЕЛЬНАЯ ЭФФЕКТИВНОСТЬ МЕЗАКОЛА И СУЛЬФАСАЛАЗИНА В ЛЕЧЕНИИ ХРОНИЧЕСКОГО РЕЦИДИВИРУЮЩЕГО ЯЗВЕННОГО КОЛИТА

Рогозина В.А., Румянцев В.Г. //Экспериментальная и клиническая гастроэнтерология, N1, 2003.

7. МЕТАБОЛИЧЕСКИЕ АСПЕКТЫ МОРФОГЕНЕЗА ЛИПИДНЫХ ВКЛЮЧЕНИЙ В ПЕЧЕНИ (ПРИГЛАШЕНИЕ К ДИСКУССИИ)

Морозов И.А. //Экспериментальная и клиническая гастроэнтерология, N1, 2003